
COVID-19 URBAN INDIAN ORGANIZATION SUMMARY JANUARY 20, 2020-SEPTEMBER 28, 2021

Albuquerque Service Area

COVID-19 Case Surveillance Information for American Indians and Alaska Natives living in counties served by Urban Indian Organizations (UIOs) in the Albuquerque Service Area

MAP 1: Albuquerque Service Area



UIO Service Area Counties	Bernalillo, Sandoval
UIO Service Area State	New Mexico



Introduction

The COVID-19 pandemic resulted in a disproportionate loss of life among American Indians and Alaska Natives. ^{1,2} To quantify this impact, Urban Indian Health Institute (UIHI) prepared this report to provide surveillance information for areas served by Urban Indian Organizations. This report contains information regarding COVID-19 cases, hospitalizations, and deaths. Please note that this data reflects information available from counties overall and does not represent individual organizations or providers.

About UIHI

As a Public Health Authority and one of 12 Tribal Epidemiology Centers in the country—and the only one that serves Urban Indian Organizations nationwide—UIHI conducts research and evaluation, collects and analyzes data, and provides disease surveillance to strengthen the health of American Indian and Alaska Native communities. UIHI's mission is to decolonize data, for Indigenous people, by Indigenous people.

Data Quality

The data used in this report is from the Centers for Disease Control and Prevention (CDC) COVID-19 Case Surveillance Data, which may differ from what is shown on your county's public health dashboard. To assess discrepancies in CDC and state data, we present metrics on data quality. The quality of COVID-19 data is measured in two ways: the completeness of racial data sent to the CDC and the proportion of cases sent to the CDC from the state.

Not all COVID-19 cases within counties have been reported to the CDC. To assess the overall representativeness of the CDC COVID-19 Case Surveillance Data, we compare the total number of COVID-19 cases reported to the CDC by county with the total number of cases reported to the New York Times.³ The number of COVID-19 cases reported by the New York Times in a given county vary based on whether they include confirmed COVID-19 cases or confirmed and probable COVID-19 cases. Therefore, the cases reported by New York Times provide an upper limit for the number of COVID-19 cases in a county.

For the data presented in this report to be of high quality, racial data needs to be both collected by states and relayed to the CDC. Unfortunately, many states have poor collection of racial data and/or do not send most COVID-19 data to the CDC. We recommend using caution when interpreting these numbers in the absence of the quality of their data. These data quality issues vary over time. Data is especially poor around the peaks of COVID-19 case burden. Please refer to Figure 1 and Table 1 to assess the overall quality of COVID-19 data in this service region.

Methods

The Centers for Disease Control and Prevention COVID-19 Case Surveillance Data are voluntarily submitted by local health jurisdictions to the CDC using the COVID-19 Case Report Form.⁴ This data is then transmitted to UIHI for analysis. Population denominators use the 2020 postcensal estimates.

The COVID-19 Case Report Form allows for the selection of multiple racial identities independent of ethnicity, however, many states and counties do not report data to the CDC in this format. Due to the methods of data collection and reporting, American Indian and Alaska Natives (Al/ANs) were defined as non-Hispanic Al/AN alone in this report. Using other definitions of Al/ANs would result in rates that underestimate the burden of COVID-19 among Al/ANs. Because of the underreporting of racial data, the statistics presented in this analysis are only applicable to non-Hispanic single race Al/ANs, a portion of the entire Al/AN population. The non-Hispanic White (NHW) population is included as a comparison population to assess disparities in COVID-19 incidence, hospitalization, and mortality in recognition of the effects of structural racism.

Ethnicity categories are captured as "Hispanic/Latino", "Non-Hispanic/Latino", or "Unknown". Age is calculated by the provided date of birth. Cases were assigned to individual counties based on their listed county of residence. Sex is recorded in the Case Report Form as "Male", "Female",

"Other", or "Unknown". The 2020 postcensal estimates do not provide estimates for individuals whose sex is not male or female, thus case rates for "Other" or "Unknown" patients cannot be calculated.

Cases were defined as individuals with a positive PCR test for COVID-19. Cases were excluded when an individual did not report county of residence. COVID-19 case dates were assigned based on when COVID-19 cases were reported to the CDC.

Age-adjusted case, hospitalization, and mortality rates stratified by race and gender are presented in this report. Age adjustment for incidence rates were calculated using the direct method applying weights from the U.S. 2000 Standard Population. Age adjustment for relative risks was performed via logistic regression.

Case counts that are less than 10 are suppressed in this report to protect confidentiality. Rates are presented with an asterisk (*) when its relative standard error (RSE) is greater than or equal to 25%. RSE provides a measure of reliability. Where the RSE is greater than or equal to 25%, the estimate is unreliable. Analyses adhere to Washington Department Of Health guidelines for the reporting of data with small numbers.⁵

Data were analyzed using R version 4.2.

GLOSSARY OF TERMS

Age-adjusted

This report refers to several analyses as age-adjusted. Age-adjustment is a statistical approach that allows for communities with different underlying age compositions to be compared. For example, COVID-19 has had disproportionate impacts based on age.⁶ As a population, Al/AN are younger than NHW.⁷ As a result of these two factors, any analysis that does not undergo age adjustment when comparing COVID-19 rates between Al/AN to NHW may only reflect that, as a population, Al/AN are younger than NHW and therefore have different rates of COVID-19. To adjust for this effect, our analysis compares individuals with similar ages to one another separately and then combines those values after appropriately weighting them.

Confirmed COVID-19

COVID-19 is determined to be "Confirmed" if there is confirmatory laboratory evidence, which requires detection of SARS-CoV-2 RNA in a clinical specimen using a molecular amplification detection test.

Confidence Interval

A confidence interval is a statistical tool used to describe the uncertainty of our estimates. It provides a range of values that is likely to include the true population value with a certain degree of confidence. It is often expressed as a percent whereby the true population value lies between an upper and lower interval. In this report, many of our estimates are presented with 95% confidence intervals. A 95% confidence interval can be interpreted as, we are 95% confident that the true value of an estimate lies within the confidence interval range.

Incidence

Incidence refers to the occurrence, rate, or frequency of disease. In this report, incidence refers to the total number of COVID-19 infections, hospitalizations, or deaths divided by the total number of individuals at risk of infection. In this report we present incidence as the number of cases per 100,000 individuals to standardize the numbers across analyses with a different total number of individuals that could have been infected with COVID-19.

Relative Risk

Relative risk in this report refers to the probability of an individual who is AI/AN being infected, hospitalized, or dying from COVID-19 relative to a non-Hispanic White individual. A relative risk greater than one indicates that AI/AN individuals had a greater risk of infection, hospitalization, or death relative to NHW. Conversely, a relative risk less than one indicates that AI/AN individuals had a lower risk relative to NHWs.

Relative Standard Error (RSE)

Standard error tells you how accurate the estimate is compared to the true value. The greater the standard error is, the more likely that the estimate is an inaccurate representation of the true value. Relative standard error is used to show how large the standard error is, relative to the size of the estimated value. For this report, we determine our estimates to be unreliable when the relative standard error of an estimate is greater than 25%. We recommend caution when using estimates with an RSE greater than 25% due to their low reliability.

Rolling Average

A rolling or moving 14-day average is the average number of confirmed COVID-19 cases, hospitalizations, or deaths over a 14-day period. It is calculated for each day by averaging the values of that day and the 13 days before. This method is used to reduce fluctuations that occur on a weekly or daily basis due to external factors such as specimen processing and reporting time.

Service Area

Urban Indian Health Organization's service areas are defined by the Urban Indian Organizations who described to UIHI the counties they serve.

EXECUTIVE SUMMARY

COVID-19 Cases

Since January 20, 2020, a total of 7,051 Al/ANs residing in the Albuquerque UIO service counties have been reported to the CDC as having confirmed COVID-19. This corresponds to an age-adjusted incidence of 13,386 COVID-19 cases per 100,000 or approximately 13 in every 100 Al/ANs in the Albuquerque Service Area were diagnosed with COVID-19. Compared to their non-Hispanic White counterparts of a similar age, Al/AN individuals were 3.13 times more likely to be infected with COVID-19.

COVID-19 Hospitalizations

Since January 20, 2020, a total of 943 Al/ANs residing in the Albuquerque UIO service counties have been reported to the CDC as hospitalized with confirmed COVID-19. This corresponds to an age-adjusted incidence of 1,846 COVID-19 hospitalizations per 100,000 people. Compared to their non-Hispanic White counterparts of similar age, Al/AN individuals were 7.11 times more likely to be hospitalized with COVID-19.

COVID-19 Deaths

Since January 20, 2020, a total of 207 Al/ANs residing in the Albuquerque UIO service counties have been reported to the CDC as having died due to COVID-19. This corresponds to an age-adjusted incidence of 217 COVID-19 deaths per 100,000 people. Compared to their non-Hispanic White counterparts of similar age, Al/AN individuals were 7.55 times more likely to die from COVID-19.

Data Quality

Of the 85,634 total COVID-19 cases reported by the New York Times in Albuquerque UIO service counties, the CDC has complete race and ethnicity available for 47,094 (55.2%) of cases. This report does not contain information regarding whether COVID-19 hospitalizations or deaths were reported to the CDC. However, of the hospitalizations and deaths in the CDC data, there is complete racial information available for 79.0% and 91.4% of cases, respectively.

FIGURE 1: Quality of COVID-19 Information Over Time, May 20, 2020–September 28, 2021, Albuquerque Service Area




TABLE 1: Overall Data Quality for Albuquerque Service Area COVID-19 Data, January 20, 2020–September 28, 2021

Data Quality Element	Data Source	Subset	# of COVID-19 Cases	Percent Completion		
Completeness of Race &	CDC	Case Reported to CDC with Complete Race & Ethnicity 47,094		55.2%		
Ethnicity in CDC Dataset	CDC	All Cases	85,278			
Completeness of Race & Ethnicity for Cases who	CDC	Case Hospitalized with Complete Race & Ethnicity	4,635	79.0%		
were Hospitalized in CDC Dataset	CDC	All Hospitalized COVID-19 Patients	All Hospitalized COVID-19 Patients 5,870			
Completeness of Race & Ethnicity for Cases who Died in CDC Dataset	CDC	COVID-19 Case Died with Complete Race & Ethnicity	1,252	91.4%		
	CDC	All COVID-19 Cases who Died	1,370			
Cases Sent to the CDC	CDC	All Cases	85,278	99.6%		
	NYT	All Cases 85,634		99.0%		
Completeness of CDC Racial Data in	CDC	Case Reported to CDC with Complete Race & Ethnicity	47,094	FF 00/		
Comparison to All Known Cases	NYT	All Cases	85,634	55.0%		

CDC = COVID-19 Case Surveillance Data Centers for Disease Control & Prevention, NYT = New York Times

TABLE 2: Incidence of COVID-19 among Non-Hispanic American Indians and Alaska Natives (Al/ANs) and Non-Hispanic Whites (NHWs), January 20, 2020– September 28, 2021, Albuquerque Service Area

	Geography	AI/AN Cases	AI/AN Population	AI/AN Age-adjusted Incidence per 100k (95% CI)	NHW Age-adjusted Incidence per 100k (95% CI)	Relative Risk Age-adjusted (95% CI)
OVERALL	UIO Service Area	7,051	46,869	13,386 (13,050, 13,732)	4,280 (4,204, 4,356)	3.13 (3.04, 3.22)
	State Overall	31,006	183,732	15,648 (15,459, 15,839)	4,916 (4,865, 4,968)	3.18 (3.13, 3.23)
BY SEX						
Female	UIO Service Area	3,852	24,928	13,579 (13,120, 14,054)	4,287 (4,182, 4,395)	3.17 (3.04, 3.30)
	State Overall	16,466	95,996	15,599 (15,342, 15,861)	5,076 (5,003, 5,151)	3.07 (3.01, 3.14)
Male	UIO Service Area	3,123	21,941	12,874 (12,389, 13,379)	4,198 (4,092, 4,307)	3.07 (2.94, 3.20)
	State Overall	14,185	87,736	15,349 (15,075, 15,628)	4,670 (4,600, 4,742)	3.29 (3.21, 3.36)
Other or Unknown	UIO Service Area	76	NA	NA	NA	NA
	State Overall	355	NA	NA	NA	NA
BY AGE						
0–19	UIO Service Area	1,421	12,611	11,268 (10,697, 11,869)	3,699 (3,531, 3,876)	3.05 (2.84, 3.27)
	State Overall	6,134	51,552	11,899 (11,605, 12,200)	3,985 (3,874, 4,099)	2.99 (2.88, 3.10)
20–54	UIO Service Area	4,274	25,023	17,080 (16,576, 17,600)	5,889 (5,759, 6,023)	2.90 (2.79, 3.01)
	State Overall	17,702	89,741	19,726 (19,437, 20,018)	6,755 (6,662, 6,849)	2.92 (2.86, 2.98)
55+	UIO Service Area	1,356	9,235	14,683 (13,922, 15,486)	3,408 (3,313, 3,505)	4.31 (4.06, 4.58)
	State Overall	7,170	42,439	16,895 (16,508, 17,290)	4,118 (4,051, 4,185)	4.10 (3.99, 4.22)
Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA

TABLE 3: Incidence of COVID-19 Hospitalizations among Non-Hispanic American Indians and Alaska Natives (AI/ANs) and Non-Hispanic Whites (NHWs), January 20, 2020 – September 28, 2021, Albuquerque Service Area

	Geography	AI/AN Cases	Al/AN Population	AI/AN Age-adjusted Incidence per 100k (95% CI)	NHW Age-adjusted Incidence per 100k (95% CI)	Relative Risk Age-adjusted (95% CI)
OVERALL	UIO Service Area	943	46,869	1,846 (1,695, 2,010)	259 (239, 282)	7.11 (6.53, 7.75)
	State Overall	4,668	183,732	1,962 (1,868, 2,061)	263 (249, 277)	7.47 (7.15, 7.80)
BY SEX						
Female	UIO Service Area	516	24,928	1,885 (1,690, 2,103)	260 (233, 290)	7.26 (6.44, 8.18)
	State Overall	2,535	95,996	1,966 (1,829, 2,115)	245 (226, 266)	8.02 (7.54, 8.53)
Male	UIO Service Area	422	21,941	1,792 (1,553, 2,068)	252 (220, 290)	7.10 (6.26, 8.05)
	State Overall	2,091	87,736	1,952 (1,824, 2,089)	279 (259, 300)	7.01 (6.58, 7.46)
Other or Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	42	NA	NA	NA	NA
BY AGE						
0–19	UIO Service Area	25	12,611	198 (134, 293)	65 (46, 92)	3.06 (1.81, 5.19)
	State Overall	93	51,552	180 (147, 221)	49 (38, 63)	3.71 (2.68, 5.14)
20–54	UIO Service Area	498	25,023	1,990 (1,823, 2,173)	267 (240, 296)	7.46 (6.51, 8.56)
	State Overall	2,196	89,741	2,447 (2,347, 2,552)	303 (284, 324)	8.06 (7.46, 8.72)
55+	UIO Service Area	420	9,235	4,548 (4,133, 5,004)	737 (694, 783)	6.17 (5.51, 6.91)
	State Overall	2,379	42,439	5,606 (5,385, 5,836)	844 (814, 875)	6.64 (6.29, 7.01)
Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA


TABLE 4: Incidence of COVID-19 Deaths among Non-Hispanic American Indians and Alaska Natives (Al/ANs) and Non-Hispanic Whites (NHWs), January 20, 2020–September 28, 2021, Albuquerque Service Area

	Geography	AI/AN Cases	Al/AN Population	Al/AN Age-adjusted Incidence per 100k (95% CI)	NHW Age-adjusted Incidence per 100k (95% CI)	Relative Risk, Age-adjusted (95% CI)
OVERALL	UIO Service Area	207	46,869	217 (156, 303)	29 (21, 40)	7.55 (6.38, 8.93)
	State Overall	1,181	183,732	263 (212, 327)	37 (30, 46)	7.06 (6.53, 7.64)
BY SEX						
Female	UIO Service Area	97	24,928	213 (148, 306)	29 (20, 42)	7.32 (5.71, 9.38)
	State Overall	529	95,996	229 (179, 291)	35 (27, 44)	6.61 (5.88, 7.43)
Male	UIO Service Area	110	21,941	279 (186, 420)	34 (22, 51)	8.28 (6.58, 10.43)
	State Overall	652	87,736	326 (241, 439)	41 (31, 56)	7.85 (7.06, 8.73)
Other or Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA
BY AGE						
0–19	UIO Service Area	<10	12,611	Suppressed	Suppressed	Suppressed
	State Overall	<10	51,552	Suppressed	Suppressed	Suppressed
20–54	UIO Service Area	52	25,023	208 (158, 273)	14 (9, 22)	14.98 (8.76, 25.60)
	State Overall	273	89,741	304 (270, 343)	25 (20, 32)	12.11 (9.37, 15.66)
55+	UIO Service Area	154	9,235	1,668 (1,424, 1,953)	310 (282, 340)	5.38 (4.48, 6.47)
	State Overall	905	42,439	2,132 (1,998, 2,276)	386 (366, 407)	5.52 (5.08, 6.01)
Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA

TABLE 5: Incidence of COVID-19 among Non-Hispanic America Indians/Alaska Natives (Al/AN) and Non-Hispanic Whites (NHW), January 20, 2020–September 28, 2021, Albuquerque Service Area


County	Geography	AI/AN Cases	Al/AN Population	Al/AN Age-adjusted Incidence per 100k (95% CI)	NHW Age-adjusted Incidence per 100k (95% CI)	Relative Risk Age-adjusted (95% CI)
Bernalillo	County	4,116	29,321	12,275 (11,877, 12,686)	4,285 (4,200, 4,371)	2.86 (2.76, 2.97)
Sandoval	County	2,935	17,548	15,161 (14,551, 15,796)	4,216 (4,051, 4,389)	3.60 (3.41, 3.80)
All New Mexico	State	31,006	183,732	15,648 (15,459, 15,839)	4,916 (4,865, 4,968)	3.18 (3.13, 3.23)


FIGURE 2: Rolling 14-day Average Incidence of COVID-19 by Date Reported to the CDC, May 20, 2020–September 28, 2021, Albuquerque Service Area



CDC COVID-19 Line Level Case Surveillance Data (Last Updated: 2021-09-28)
Denominators from: 2020 Postcensal Estimates


FIGURE 3: Rolling 14-day Average Incidence of COVID-19 Hospitalizations by Date Reported to the CDC, May 20, 2020– September 28, 2021, Albuquerque Service Area





CDC COVID-19 Line Level Case Surveillance Data (Last Updated: 2021-09-28)
Denominators from: 2020 Postcensal Estimates





CDC COVID-19 Line Level Case Surveillance Data (Last Updated: 2021-09-28)
Denominators from: 2020 Postcensal Estimates

To request technical assistance in interpreting this report, or to request additional information, email Scott Erickson at scotte@uihi.org

Bibliography:

- 1. Arrazola J, Masiello MM, Joshi S, et al. COVID-19 Mortality Among American Indian and Alaska Native Persons 14 States, January-June 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(49):1853-1856. doi:10.15585/mmwr.mm6949a3
- Sarah M. Hatcher et al., COVID-19 Among American Indian and Alaska Native Persons 23 States, January 31–July 3, 2020. MMWR Morb Mortal Wkly Rep. 2020;69. doi:10.15585/mmwr.mm6934e1
- 3. The New York Times. Coronavirus (Covid-19) Data in the United States. Published online 2021. Accessed June 18, 2021. https://github.com/nytimes/covid-19-data
- Centers for Disease Control & Prevention. Human Infection with 2019 Novel Coronavirus Case Report Form. Published online January 15, 2021. Accessed June 18, 2021. https://www.cdc.gov/coronavirus/2019-ncov/downloads/pui-form.pdf
- Washington Department of Health. Department of Health Agency Standards for Reporting Data with Small Numbers. Published online May 2018. Accessed June 18, 2021. https://www.doh.wa.gov/Portals/1/Documents/1500/SmallNumbers.pdf
- Center for Disease Control and Prevention. Demographic Trends of COVID-19 cases and deaths in the US reported to CDC. Published online July 22, 2021. https://covid.cdc.gov/covid-data-tracker/#demographics
- 7. Urban Indian Health Institute. Community Health Profile: National Aggregate of Urban Indian Program Service Areas.; 2016:58.