
COVID-19 URBAN INDIAN ORGANIZATION SUMMARY JANUARY 20, 2020-SEPTEMBER 28, 2021

Salt Lake City Service Area

COVID-19 Case Surveillance Information for American Indians and Alaska Natives living in counties served by Urban Indian Organizations (UIOs) in the Salt Lake City Service Area

MAP 1: Salt Lake City Service Area



UIO Service Area Counties	Davis, Salt Lake, Tooele, Utah, Weber
UIO Service Area State	Utah



Introduction

The COVID-19 pandemic resulted in a disproportionate loss of life among American Indians and Alaska Natives. ^{1,2} To quantify this impact, Urban Indian Health Institute (UIHI) prepared this report to provide surveillance information for areas served by Urban Indian Organizations. This report contains information regarding COVID-19 cases, hospitalizations, and deaths. Please note that this data reflects information available from counties overall and does not represent individual organizations or providers.

About UIHI

As a Public Health Authority and one of 12 Tribal Epidemiology Centers in the country—and the only one that serves Urban Indian Organizations nationwide—UIHI conducts research and evaluation, collects and analyzes data, and provides disease surveillance to strengthen the health of American Indian and Alaska Native communities. UIHI's mission is to decolonize data, for Indigenous people, by Indigenous people.

Data Quality

The data used in this report is from the Centers for Disease Control and Prevention (CDC) COVID-19 Case Surveillance Data, which may differ from what is shown on your county's public health dashboard. To assess discrepancies in CDC and state data, we present metrics on data quality. The quality of COVID-19 data is measured in two ways: the completeness of racial data sent to the CDC and the proportion of cases sent to the CDC from the state.

Not all COVID-19 cases within counties have been reported to the CDC. To assess the overall representativeness of the CDC COVID-19 Case Surveillance Data, we compare the total number of COVID-19 cases reported to the CDC by county with the total number of cases reported to the New York Times.³ The number of COVID-19 cases reported by the New York Times in a given county vary based on whether they include confirmed COVID-19 cases or confirmed and probable COVID-19 cases. Therefore, the cases reported by New York Times provide an upper limit for the number of COVID-19 cases in a county.

For the data presented in this report to be of high quality, racial data needs to be both collected by states and relayed to the CDC. **Unfortunately, many states have poor collection of racial data and/or do not send most COVID-19 data to the CDC. We recommend using caution when interpreting these numbers in the absence of the quality of their data.** These data quality issues vary over time. Data is especially poor around the peaks of COVID-19 case burden. Please refer to Figure 1 and Table 1 to assess the overall quality of COVID-19 data in this service region.

Methods

The Centers for Disease Control and Prevention COVID-19 Case Surveillance Data are voluntarily submitted by local health jurisdictions to the CDC using the COVID-19 Case Report Form.⁴ This data is then transmitted to UIHI for analysis. Population denominators use the 2020 postcensal estimates.

The COVID-19 Case Report Form allows for the selection of multiple racial identities independent of ethnicity, however, many states and counties do not report data to the CDC in this format. Due to the methods of data collection and reporting, American Indian and Alaska Natives (Al/ANs) were defined as non-Hispanic Al/AN alone in this report. Using other definitions of Al/ANs would result in rates that underestimate the burden of COVID-19 among Al/ANs. Because of the underreporting of racial data, the statistics presented in this analysis are only applicable to non-Hispanic single race Al/ANs, a portion of the entire Al/AN population. The non-Hispanic White (NHW) population is included as a comparison population to assess disparities in COVID-19 incidence, hospitalization, and mortality in recognition of the effects of structural racism.

Ethnicity categories are captured as "Hispanic/Latino", "Non-Hispanic/Latino", or "Unknown". Age is calculated by the provided date of birth. Cases were assigned to individual counties based on their listed county of residence. Sex is recorded in the Case Report Form as "Male", "Female",

"Other", or "Unknown". The 2020 postcensal estimates do not provide estimates for individuals whose sex is not male or female, thus case rates for "Other" or "Unknown" patients cannot be calculated.

Cases were defined as individuals with a positive PCR test for COVID-19. Cases were excluded when an individual did not report county of residence. COVID-19 case dates were assigned based on when COVID-19 cases were reported to the CDC.

Age-adjusted case, hospitalization, and mortality rates stratified by race and gender are presented in this report. Age adjustment for incidence rates were calculated using the direct method applying weights from the U.S. 2000 Standard Population. Age adjustment for relative risks was performed via logistic regression.

Case counts that are less than 10 are suppressed in this report to protect confidentiality. Rates are presented with an asterisk (*) when its relative standard error (RSE) is greater than or equal to 25%. RSE provides a measure of reliability. Where the RSE is greater than or equal to 25%, the estimate is unreliable. Analyses adhere to Washington Department Of Health guidelines for the reporting of data with small numbers.⁵

Data were analyzed using R version 4.2.

GLOSSARY OF TERMS

Age-adjusted

This report refers to several analyses as age-adjusted. Age-adjustment is a statistical approach that allows for communities with different underlying age compositions to be compared. For example, COVID-19 has had disproportionate impacts based on age.⁶ As a population, Al/AN are younger than NHW.⁷ As a result of these two factors, any analysis that does not undergo age adjustment when comparing COVID-19 rates between Al/AN to NHW may only reflect that, as a population, Al/AN are younger than NHW and therefore have different rates of COVID-19. To adjust for this effect, our analysis compares individuals with similar ages to one another separately and then combines those values after appropriately weighting them.

Confirmed COVID-19

COVID-19 is determined to be "Confirmed" if there is confirmatory laboratory evidence, which requires detection of SARS-CoV-2 RNA in a clinical specimen using a molecular amplification detection test.

Confidence Interval

A confidence interval is a statistical tool used to describe the uncertainty of our estimates. It provides a range of values that is likely to include the true population value with a certain degree of confidence. It is often expressed as a percent whereby the true population value lies between an upper and lower interval. In this report, many of our estimates are presented with 95% confidence intervals. A 95% confidence interval can be interpreted as, we are 95% confident that the true value of an estimate lies within the confidence interval range.

Incidence

Incidence refers to the occurrence, rate, or frequency of disease. In this report, incidence refers to the total number of COVID-19 infections, hospitalizations, or deaths divided by the total number of individuals at risk of infection. In this report we present incidence as the number of cases per 100,000 individuals to standardize the numbers across analyses with a different total number of individuals that could have been infected with COVID-19.

Relative Risk

Relative risk in this report refers to the probability of an individual who is AI/AN being infected, hospitalized, or dying from COVID-19 relative to a non-Hispanic White individual. A relative risk greater than one indicates that AI/AN individuals had a greater risk of infection, hospitalization, or death relative to NHW. Conversely, a relative risk less than one indicates that AI/AN individuals had a lower risk relative to NHWs.

Relative Standard Error (RSE)

Standard error tells you how accurate the estimate is compared to the true value. The greater the standard error is, the more likely that the estimate is an inaccurate representation of the true value. Relative standard error is used to show how large the standard error is, relative to the size of the estimated value. For this report, we determine our estimates to be unreliable when the relative standard error of an estimate is greater than 25%. We recommend caution when using estimates with an RSE greater than 25% due to their low reliability.

Rolling Average

A rolling or moving 14-day average is the average number of confirmed COVID-19 cases, hospitalizations, or deaths over a 14-day period. It is calculated for each day by averaging the values of that day and the 13 days before. This method is used to reduce fluctuations that occur on a weekly or daily basis due to external factors such as specimen processing and reporting time.

Service Area

Urban Indian Organization's service areas are defined by the Urban Indian Organizations who described to UIHI the counties they serve.

EXECUTIVE SUMMARY

COVID-19 Cases

Since January 20, 2020, a total of 2,309 Al/ANs residing in the Salt Lake City UIO service counties have been reported to the CDC as having confirmed COVID-19. This corresponds to an age-adjusted incidence of 13,604 COVID-19 cases per 100,000 or approximately 13 in every 100 Al/ANs in the Salt Lake City Service Area were diagnosed with COVID-19. Compared to their non-Hispanic White counterparts of a similar age, Al/AN individuals were 1.22 times more likely to be infected with COVID-19.

COVID-19 Hospitalizations

Since January 20, 2020, a total of 111 Al/ANs residing in the Salt Lake City UIO service counties have been reported to the CDC as hospitalized with confirmed COVID-19. This corresponds to an age-adjusted incidence of 695 COVID-19 hospitalizations per 100,000 people. Compared to their non-Hispanic White counterparts of similar age, Al/AN individuals were 4.52 times more likely to be hospitalized with COVID-19.

COVID-19 Deaths

Since January 20, 2020, a total of 29 Al/ANs residing in the Salt Lake City UIO service counties have been reported to the CDC as having died due to COVID-19. This corresponds to an age-adjusted incidence of 102 COVID-19 deaths per 100,000 people. Compared to their non-Hispanic White counterparts of similar age, Al/AN individuals were 5.24 times more likely to die from COVID-19.

Data Quality

Of the 397,646 total COVID-19 cases reported by the New York Times in Salt Lake City UIO service counties, the CDC has complete race and ethnicity available for 317,510 (79.8%) of cases. This report does not contain information regarding whether COVID-19 hospitalizations or deaths were reported to the CDC. However, of the hospitalizations and deaths in the CDC data, there is complete racial information available for 87.8% and 85.8% of cases, respectively.

FIGURE 1: Quality of COVID-19 Information Over Time, May 20, 2020–September 28, 2021, Salt Lake City Service Area




TABLE 1: Overall Data Quality for Salt Lake City Service Area COVID-19 Data, January 20, 2020–September 28, 2021

Data Quality Element	Data Source	Subset	# of COVID-19 Cases	Percent Completion	
Completeness of Race & Ethnicity in CDC Dataset	CDC	Case Reported to CDC with Complete Race & Ethnicity	317,510	- 82.2%	
	CDC	All Cases	386,407	<i>32.2 /</i> 0	
Completeness of Race & Ethnicity for Cases who were Hospitalized in CDC Dataset	CDC	Case Hospitalized with Complete Race & Ethnicity	6,521	87.8%	
	CDC	All Hospitalized COVID-19 Patients	7,429		
Completeness of Race & Ethnicity for Cases who Died in CDC Dataset	CDC	COVID-19 Case Died with Complete Race & Ethnicity	1,782	85.8%	
	CDC	All COVID-19 Cases who Died	2,077	00.076	
Cases Sent to the CDC	CDC	All Cases	386,407	- 97.2%	
	NYT	All Cases	397,646		
Completeness of CDC Racial Data in	CDC	Case Reported to CDC with Complete Race & Ethnicity	317,510	- 79.8%	
Comparison to All Known Cases	NYT	All Cases	397,646	10.070	

CDC = COVID-19 Case Surveillance Data Centers for Disease Control & Prevention, NYT = New York Times

TABLE 2: Incidence of COVID-19 among Non-Hispanic American Indians and Alaska Natives (Al/ANs) and Non-Hispanic Whites (NHWs), January 20, 2020– September 28, 2021, Salt Lake City Service Area

	Geography	AI/AN Cases	Al/AN Population	Al/AN Age-adjusted Incidence per 100k (95% CI)	NHW Age-adjusted Incidence per 100k (95% CI)	Relative Risk Age-adjusted (95% CI)
OVERALL	UIO Service Area	2,309	14,107	13,604 (13,057, 14,173)	11,117 (11,057, 11,177)	1.22 (1.17, 1.27)
	State Overall	4,745	30,125	13,388 (13,010, 13,777)	10,923 (10,873, 10,974)	1.23 (1.19, 1.26)
BY SEX						
Female	UIO Service Area	1,239	7,125	14,180 (13,408, 14,997)	11,247 (11,163, 11,331)	1.26 (1.19, 1.33)
	State Overall	2,642	15,332	14,399 (13,856, 14,963)	11,058 (10,987, 11,129)	1.30 (1.25, 1.35)
Male	UIO Service Area	1,069	6,982	12,996 (12,235, 13,803)	10,992 (10,906, 11,079)	1.18 (1.11, 1.26)
	State Overall	2,100	14,793	12,300 (11,782, 12,841)	10,788 (10,716, 10,860)	1.14 (1.09, 1.19)
Other or Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA
BY AGE						
0–19	UIO Service Area	481	3,827	12,569 (11,494, 13,744)	9,521 (9,441, 9,602)	1.32 (1.21, 1.44)
	State Overall	1,058	8,878	11,917 (11,220, 12,657)	9,250 (9,181, 9,319)	1.29 (1.21, 1.37)
20–54	UIO Service Area	1,494	8,105	18,433 (17,522, 19,392)	15,894 (15,813, 15,976)	1.16 (1.10, 1.22)
	State Overall	2,821	15,529	18,166 (17,508, 18,849)	15,761 (15,690, 15,833)	1.15 (1.11, 1.20)
55+	UIO Service Area	334	2,175	15,356 (13,795, 17,095)	10,185 (10,089, 10,282)	1.51 (1.35, 1.68)
	State Overall	866	5,718	15,145 (14,169, 16,188)	10,019 (9,939, 10,099)	1.51 (1.41, 1.62)
Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA

TABLE 3: Incidence of COVID-19 Hospitalizations among Non-Hispanic American Indians and Alaska Natives (Al/ANs) and Non-Hispanic Whites (NHWs), January 20, 2020 – September 28, 2021, Salt Lake City Service Area

	Geography	AI/AN Cases	Al/AN Population	Al/AN Age-adjusted Incidence per 100k (95% CI)	NHW Age-adjusted Incidence per 100k (95% CI)	Relative Risk Age-adjusted (95% CI)
OVERALL	UIO Service Area	111	14,107	695 (573, 843)	154 (145, 163)	4.52 (3.74, 5.46)
	State Overall	215	30,125	562 (487, 648)	130 (123, 138)	4.31 (3.76, 4.94)
BY SEX						
Female	UIO Service Area	64	7,125	767 (593, 992)	139 (127, 153)	5.50 (4.28, 7.06)
	State Overall	114	15,332	571 (468, 697)	119 (109, 130)	4.80 (3.98, 5.80)
Male	UIO Service Area	47	6,982	623 (465, 836)	170 (157, 184)	3.67 (2.75, 4.90)
	State Overall	101	14,793	560 (456, 687)	143 (132, 154)	3.92 (3.22, 4.79)
Other or Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA
BY AGE						
0–19	UIO Service Area	<10	3,827	Suppressed	14 (11, 18)	Suppressed
	State Overall	<10	8,878	Suppressed	11 (9, 14)	Suppressed
20–54	UIO Service Area	66	8,105	814 (640, 1,036)	147 (140, 155)	5.53 (4.32, 7.08)
	State Overall	104	15,529	670 (553, 812)	128 (122, 135)	5.23 (4.28, 6.38)
55+	UIO Service Area	43	2,175	1,977 (1,466, 2,666)	664 (639, 689)	2.98 (2.20, 4.03)
	State Overall	106	5,718	1,854 (1,532, 2,243)	565 (546, 584)	3.28 (2.71, 3.98)
Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA

TABLE 4: Incidence of COVID-19 Deaths among Non-Hispanic American Indians and Alaska Natives (Al/ANs) and Non-Hispanic Whites (NHWs), January 20, 2020–September 28, 2021, Salt Lake City Service Area

	Geography	AI/AN Cases	Al/AN Population	Al/AN Age-adjusted Incidence per 100k (95% CI)	NHW Age-adjusted Incidence per 100k (95% CI)	Relative Risk, Age-adjusted (95% CI)
OVERALL	UIO Service Area	29	14,107	102 (65, 160)	19 (15, 26)	5.24 (3.62, 7.58)
	State Overall	79	30,125	98 (70, 137)	19 (15, 25)	5.12 (4.08, 6.41)
BY SEX						
Female	UIO Service Area	14	7,125	95 (52, 174)*	17 (12, 24)	5.48 (3.22, 9.35)*
	State Overall	32	15,332	80 (51, 126)	17 (13, 24)	4.63 (3.25, 6.60)
Male	UIO Service Area	15	6,982	133 (75, 237)*	26 (20, 35)	5.07 (3.03, 8.48)*
	State Overall	47	14,793	143 (97, 211)	25 (19, 33)	5.66 (4.22, 7.59)
Other or Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA
BY AGE						
0–19	UIO Service Area	<10	3,827	Suppressed	Suppressed	Suppressed
	State Overall	<10	8,878	Suppressed	Suppressed	Suppressed
20–54	UIO Service Area	<10	8,105	Suppressed	14 (12, 16)	Suppressed
	State Overall	20	15,529	129 (83, 200)	15 (13, 18)	8.46 (5.33, 13.42)
55+	UIO Service Area	22	2,175	1,011 (666, 1,536)	297 (281, 314)	3.41 (2.24, 5.20)
	State Overall	59	5,718	1,032 (799, 1,332)	290 (277, 304)	3.56 (2.75, 4.61)
Unknown	UIO Service Area	<10	NA	NA	NA	NA
	State Overall	<10	NA	NA	NA	NA


^{*=}Relative Standard Error > 25%, Estimate Uncertain

TABLE 5: Incidence of COVID-19 among Non-Hispanic America Indians/Alaska Natives (AI/AN) and Non-Hispanic Whites (NHW), January 20, 2020–September 28, 2021, Salt Lake City Service Area


County	Geography	AI/AN Cases	Al/AN Population	Al/AN Age-adjusted Incidence per 100k (95% CI)	NHW Age-adjusted Incidence per 100k (95% CI)	Relative Risk Age-adjusted (95% CI)
Davis	County	249	1,583	13,530 (11,945, 15,326)	10,975 (10,824, 11,128)	1.23 (1.09, 1.40)*
Salt Lake	County	1,283	7,663	13,862 (13,119, 14,648)	10,676 (10,589, 10,763)	1.30 (1.23, 1.37)
Tooele	County	76	567	10,886 (8,680, 13,653)	8,716 (8,390, 9,055)	1.25 (1.00, 1.57)*
Utah	County	498	2,910	14,385 (13,171, 15,712)	12,905 (12,773, 13,039)	1.11 (1.02, 1.22)*
Weber	County	203	1,384	12,183 (10,612, 13,986)	10,274 (10,103, 10,448)	1.19 (1.03, 1.36)*
All Utah	State	2,309	14,107	13,604 (13,057, 14,173)	11,117 (11,057, 11,177)	1.22 (1.17, 1.27)


^{*=}Relative Standard Error > 25%, Estimate Uncertain

FIGURE 2: Rolling 14-day Average Incidence of COVID-19 by Date Reported to the CDC, May 20, 2020–September 28, 2021, Salt Lake City Service Area




CDC COVID-19 Line Level Case Surveillance Data (Last Updated: 2021-09-28)
Denominators from: 2020 Postcensal Estimates





CDC COVID-19 Line Level Case Surveillance Data (Last Updated: 2021-09-28)
Denominators from: 2020 Postcensal Estimates





CDC COVID-19 Line Level Case Surveillance Data (Last Updated: 2021-09-28)
Denominators from: 2020 Postcensal Estimates

To request technical assistance in interpreting this report, or to request additional information, email Scott Erickson at scotte@uihi.org

Bibliography:

- 1. Arrazola J, Masiello MM, Joshi S, et al. COVID-19 Mortality Among American Indian and Alaska Native Persons 14 States, January-June 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(49):1853-1856. doi:10.15585/mmwr.mm6949a3
- Sarah M. Hatcher et al., COVID-19 Among American Indian and Alaska Native Persons 23 States, January 31–July 3, 2020. MMWR Morb Mortal Wkly Rep. 2020;69. doi:10.15585/mmwr.mm6934e1
- 3. The New York Times. Coronavirus (Covid-19) Data in the United States. Published online 2021. Accessed June 18, 2021. https://github.com/nytimes/covid-19-data
- 4. Centers for Disease Control & Prevention. Human Infection with 2019 Novel Coronavirus Case Report Form. Published online January 15, 2021. Accessed June 18, 2021. https://www.cdc.gov/coronavirus/2019-ncov/downloads/pui-form.pdf
- Washington Department of Health. Department of Health Agency Standards for Reporting Data with Small Numbers. Published online May 2018. Accessed June 18, 2021. https://www.doh.wa.gov/Portals/1/Documents/1500/SmallNumbers.pdf
- Center for Disease Control and Prevention. Demographic Trends of COVID-19 cases and deaths in the US reported to CDC. Published online July 22, 2021. https://covid.cdc.gov/covid-data-tracker/#demographics
- 7. Urban Indian Health Institute. Community Health Profile: National Aggregate of Urban Indian Health Program Service Areas.; 2016:58.